

APPENDIX E
RANGELAND PRODUCTIVITY AND PLANT COMPOSITION

Rangeland Productivity and Plant Composition

Comanche County, Oklahoma

[Only the soils that support rangeland vegetation suitable for grazing are rated. This report shows only the major soils in each map unit]

Map symbol and soil name	Ecological site	Total dry-weight production			Characteristic vegetation	Rangeland composition
		Favorable year	Normal year	Unfavorable year		
		Lb/ac	Lb/ac	Lb/ac		Pct
VeC: Vernon	Clay prairie (South) PE 31-44	2,500	1,700	1,000	Sideoats grama	25
					Buffalograss	15
					Tobosa	15
					Other perennial grasses	10
					Blue grama	5
					Hairy grama	5
					Other annual forbs	5
					Other perennial forbs	5
					Other shrubs	5
					Silver bluestem	5
					Vine mesquite	5

Rangeland Productivity and Plant Composition

In areas that have similar climate and topography, differences in the kind and amount of rangeland or forest understory vegetation are closely related to the kind of soil. Effective management is based on the relationship between the soils and vegetation and water.

This table shows, for each soil that supports vegetation suitable for grazing, the ecological site; the total annual production of vegetation in favorable, normal, and unfavorable years; the characteristic vegetation; and the average percentage of each species. An explanation of the column headings in the table follows.

An "ecological site" is the product of all the environmental factors responsible for its development. It has characteristic soils that have developed over time throughout the soil development process; a characteristic hydrology, particularly infiltration and runoff that has developed over time; and a characteristic plant community (kind and amount of vegetation). The hydrology of the site is influenced by development of the soil and plant community. The vegetation, soils, and hydrology are all interrelated. Each is influenced by the others and influences the development of the others. The plant community on an ecological site is typified by an association of species that differs from that of other ecological sites in the kind and/or proportion of species or in total production. Descriptions of ecological sites are provided in the Field Office Technical Guide, which is available in local offices of the Natural Resources Conservation Service (NRCS).

"Total dry-weight production" is the amount of vegetation that can be expected to grow annually in a well managed area that is supporting the potential natural plant community. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruits of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation for favorable, normal, and unfavorable years. In a favorable year, the amount and distribution of precipitation and the temperatures make growing conditions substantially better than average. In a normal year, growing conditions are about average. In an unfavorable year, growing conditions are well below average, generally because of low available soil moisture. Yields are adjusted to a common percent of air-dry moisture content.

"Characteristic vegetation" (the grasses, forbs, and shrubs that make up most of the potential natural plant community on each soil) is listed by common name. Under "rangeland composition," the expected percentage of the total annual production is given for each species making up the characteristic vegetation. The amount that can be used as forage depends on the kinds of grazing animals and on the grazing season.

Range management requires knowledge of the kinds of soil and of the potential natural plant community. It also requires an evaluation of the present range similarity index and rangeland trend. Range similarity index is determined by comparing the present plant community with the potential natural plant community on a particular rangeland ecological site. The more closely the existing community resembles the potential community, the higher the range similarity index. Rangeland trend is defined as the direction of change in an existing plant community relative to the potential natural plant community. Further information about the range similarity index and rangeland trend is available in the "National Range and Pasture Handbook," which is available in local offices of NRCS or on the Internet.

The objective in range management is to control grazing so that the plants growing on a site are about the same in kind and amount as the potential natural plant community for that site. Such management generally results in the optimum production of vegetation, control of undesirable brush species, conservation of water, and control of erosion. Sometimes, however, an area with a range similarity index somewhat below the potential meets grazing needs, provides wildlife habitat, and protects soil and water resources.

Reference:

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. (<http://www.nrcs.usda.gov/technical/range.html>)

Rangeland Productivity and Plant Composition

Comanche County, Oklahoma

[Only the soils that support rangeland vegetation suitable for grazing are rated. This report shows only the major soils in each map unit]

Map symbol and soil name	Ecological site	Total dry-weight production			Characteristic vegetation	Rangeland composition
		Favorable year	Normal year	Unfavorable year		
		<i>Lb/ac</i>	<i>Lb/ac</i>	<i>Lb/ac</i>		<i>Pct</i>
Pc: Ashport	Loamy bottomland PE 44-64	8,500	6,100	4,500	Big bluestem	25
					Indiangrass	15
					Other perennial grasses	15
					Switchgrass	15
					Little bluestem	10
					Other perennial forbs	10
					Eastern gamagrass	5
					Other trees	5